COLONOSCOPY – DIET / LAXATIVE BOWEL PREPARATION – MULTIDAY SUPREP

Hillmont GI. P.C. 215.402.0800

WHAT TO EXPECT:

- Most people have a bowel movement within an hour or two of starting the laxative. Sometimes, there may be a delay of several hours. Be patient and stay close to a bathroom. Multiple bowel movements will occur and may continue for several hours after you have finished drinking the solution
- Your stools should become clear of solid material. When you are fully prepared the final bowel movements should look like urine in the toilet water
- Bloating, chills and/or nausea are common after the first few glasses and more common if the liquids are chilled. This is
 usually temporary and improves once bowel movements begin. If you develop vomiting, severe discomfort or bloating,
 stop taking the prep for a while until the symptoms go away. If vomiting persists, stop the preparation and call our office
 215-402-0800

7 DAYS PRIOR TO YOUR PROCEDURE: Review Medication handout, obtain laxative preparation 3 DAYS PRIOR TO YOUR PROCEDURE: AVOID WHOLE GRAINS, NUTS, CORN, POPCORN & SMALL SEEDS

- Whole grains- oatmeal, granola, 7 grain breads, rice, etc.
- Seeds raspberries, strawberries, cucumbers, tomatoes, poppy, sesame, seeds, etc

_____ DAYS BEFORE PROCEDURE – Purchase Miralax in the laxative aisle and start taking a serving (one capful/17 gm/heaping tbsp in at least 8 ounces of water) once / twice daily.

_____TWO DAYS BEFORE PROCEDURE – Purchase Citrate of Magnesia in the laxative aisle. At night, drink entire bottle (10 ounces) and at least two glasses of water. It usually works between ½ and 4 hours.

THE DAY BEFORE THE PROCEDURE: NO SOLID FOOD AT ALL. CLEAR LIQUIDS ALL DAY

- NO CLOUDY LIQUIDS, milk, milk drinks, creamer, liquids that have pulp, NO alcoholic beverages CLEAR LIQUIDS: include any liquid that you can see clearly through. Avoid RED and PURPLE
- Water
- Carbonated beverages / soda
- Gatorade® Kool-aid®
- Jello® Popsicles or water ice
- Coffee or Tea- without milk, non-dairy creamer or cream
- Clear fruit juices without pulp apple, grape, lemonade, etc
- Clear hard candy, sugar, and lemon
- Clear broth (chicken, beef, vegetable, etc)

IMPORTANT NOTE: FOLLOW THESE INSTRUCTIONS AND NOT the instructions on the Prep's packaging. THE NIGHT BEFORE YOUR PROCEDURE: 7:00 pm

- Pour ONE (1) 6-ounce bottle of SUPREP liquid into the mixing container. Add water to the 16-ounce line on the container and mix. Take your time and slowly finish the dose within 30 minutes. Drink ALL of the solution
- You MUST drink two (2) more 16-ounce containers of water over the next hour
- Drink at least an additional 16-ounces of water or clear liquid before bedtime. Drinking more usually leads to a better
 preparation and complete and thorough examination

THE DAY OF YOUR PROCEDURE: FIVE (5) hours before your procedure, AT _____ AM/PM

- Pour the 2nd 6-ounce bottle of SUPREP liquid into the mixing container. Add water to the 16-ounce line on the container and mix. Take your time and slowly finish the dose within 30 minutes. **Drink ALL of the solution**
- Drink TWO more 16-ounce containers of water over the next hour
- Drink at least an additional 16-ounces of water or clear liquid before you need to stop drinking
- You MUST stop drinking 3 hours before your procedure time, so nothing after
- NO SMOKING the day of the procedure
- You must put NOTHING in your mouth for the last 3 hours prior to your procedure. This means NO water, ice chips, gum, candy, cough drops, etc., except that you should take your medications with a sip of water

For questions about the preparation, medications or to reschedule your appointment, call 215-402-0800 between 9:00AM and 5:00PM