## SPECIAL INSTRUCTION FOR MEDICATIONS IN PREPARATION FOR PROCEDURE

Hillmont GI, P.C. 215 402.0800

NIANAE /DOD.

NAIVIE/DUB:
STARTS 1 WEEK PRIOR TO PROCEDURE
To have a complete and accurate and examination done as safely as possible, it is important that you follow the medication recommendations on this page. If these are not followed, the test may have to be rescheduled for another day.
SPECIAL INSTRUCTIONS
☐ BLOOD THINNERS
STOP blood thinner DAYS prior to the procedure
☐ DIABETIC MEDICATIONS
DAY BEFORE PROCEDURE: NO CHANGE TAKE NONE ALL DAY SKIP EVENING MEDS  OR
DAY OF THE PROCEDURE:   NO CHANGE   NONE UNTIL YOU RESUME DIET  OR
NO DIURETICS/WATER PILLS ON THE DAY OF THE PROCEDURE Includes: Lasix(Furosemide), Bumex(Bumetanide), Dyazide(HCTZ/Hydrochlorothiazide), Demadex(Torsemide), and others
☐ OTHER:
OTHERWISE, TAKE ALL MEDICATIONS AS YOU NORMALLY WOULD. IF YOU NEED TO TAKE MEDICATIONS WITHIN 3 HOURS OF THE PROCEDURE, THEN DO IT WITH ONLY A SIP OF WATER.  For questions about the procedure, medications or to reschedule your appointment, call 215-402-0800 between 9:00AM and 5:00PM
CLINICIAN: